

(6) 琵琶湖南湖年次別富栄養化項目平均値(昭和54年度～平成17年度)

水域	項目	全窒素	NO ₂ ⁻ -N	NO ₃ ⁻ -N	有機態-N	全りん	PO ₄ ³⁻	クロロフィルa
		mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	μg/L
南	54	0.41	0.003	0.08	0.30	0.034	0.011	13.5
	55	0.41	0.005	0.09	0.28	0.027	0.012	11.7
	56	0.42	0.004	0.09	0.30	0.022	0.007	12.8
	57	0.40	0.004	0.09	0.28	0.025	0.006	11.1
	58	0.35	0.003	0.10	0.22	0.021	0.005	10.0
	59	0.37	0.004	0.12	0.22	0.022	0.007	7.3
	60	0.41	0.003	0.11	0.28	0.027	0.009	11.8
	61	0.37	0.004	0.10	0.24	0.024	0.006	9.3
	62	0.34	0.003	0.07	0.26	0.022	0.007	9.5
	63	0.41	0.004	0.13	0.26	0.024	0.008	10.1
湖	1	0.39	0.004	0.12	0.24	0.022	0.003	9.4
	2	0.40	0.004	0.14	0.24	0.025	0.005	9.8
	3	0.39	0.004	0.13	0.24	0.023	0.003	9.4
	4	0.41	0.004	0.10	0.29	0.024	0.003	12.5
	5	0.39	0.004	0.12	0.25	0.020	0.002	10.6
	6	0.39	0.003	0.11	0.26	0.022	0.003	8.1
	7	0.44	0.004	0.17	0.25	0.020	0.003	9.2
	8	0.42	0.004	0.16	0.25	0.018	0.003	7.9
	9	0.42	0.004	0.17	0.23	0.019	0.004	7.0
	10	0.40	0.004	0.15	0.24	0.018	0.004	7.2
	11	0.39	0.004	0.13	0.24	0.018	0.004	8.2
	12	0.40	0.003	0.14	0.24	0.019	0.006	5.6
	13	0.32	0.003	0.11	0.20	0.019	0.006	6.0
	14	0.34	0.003	0.12	0.20	0.018	0.007	6.7
15	0.36	0.003	0.15	0.20	0.016	0.005	5.8	
16	0.33	0.002	0.13	0.20	0.016	0.007	5.0	
17	0.32	0.002	0.11	0.20	0.017	0.009	5.8	