

原発部位	5年予後 診断年	全症例 (上皮内がんを除く)(初発のみ)					手術(原発巣切除)あり					手術(原発巣切除)なし (姑息・対症療法ありを含む)				
		生存	死亡	不明	計	予後 判明率	生存	死亡	不明	計	予後 判明率	生存	死亡	不明	計	予後 判明率
全部位	2005	374	375	50	799	93.7%	307	161	40	508	92.1%	67	214	10	291	96.6%
	2006	432	387	16	835	98.1%	358	158	12	528	97.7%	74	229	4	307	98.7%
	2007	413	307	3	723	99.6%	344	125	1	470	99.8%	69	182	2	253	99.2%
	2008	437	331	12	780	98.5%	362	113	10	485	97.9%	75	218	2	295	99.3%
	2009	425	353	17	795	97.9%	356	159	11	526	97.9%	69	194	6	269	97.8%
	2010	431	409	13	853	98.5%	371	172	7	550	98.7%	60	237	6	303	98.0%
	2011	493	441	33	967	96.6%	386	170	15	571	97.4%	107	271	18	396	95.5%
	2012	498	430	55	983	94.4%	391	144	29	564	94.9%	107	286	26	419	93.8%
	2013	462	362	83	907	90.8%	386	171	57	614	90.7%	76	191	26	293	91.1%
	2014	333	405	275	1,013	72.9%	271	158	217	646	66.4%	62	247	58	367	84.2%
2005~2014	4,298	3,800	557	8,655	93.6%	3,532	1,531	399	5,462	92.7%	766	2,269	158	3,193	95.1%	
食道	2005	4	18	2	24	91.7%	3	8	1	12	91.7%	1	10	1	12	91.7%
	2006	5	24		29	100.0%	4	13		17	100.0%	1	11		12	100.0%
	2007	4	12		16	100.0%	4	3		7	100.0%		9		9	100.0%
	2008	9	14		23	100.0%	6	5		11	100.0%	3	9		12	100.0%
	2009	4	19		23	100.0%	4	6		10	100.0%		13		13	100.0%
	2010	8	21		29	100.0%	5	9		14	100.0%	3	12		15	100.0%
	2011	9	20		29	100.0%	7	6		13	100.0%	2	14		16	100.0%
	2012	8	14	2	24	91.7%	6	7		13	100.0%	2	7	2	11	81.8%
	2013	6	16	1	23	95.7%	3	10	1	14	92.9%	3	6		9	100.0%
	2014	7	15	8	30	73.3%	4	5	5	14	64.3%	3	10	3	16	81.3%
2005~2014	64	173	13	250	94.8%	46	72	7	125	94.4%	18	101	6	125	95.2%	
胃	2005	76	67	20	163	87.7%	75	48	20	143	86.0%	1	19		20	100.0%
	2006	89	66	3	158	98.1%	88	41	3	132	97.7%	1	25		26	100.0%
	2007	81	48	1	130	99.2%	78	29	1	108	99.1%	3	19		22	100.0%
	2008	75	52	5	132	96.2%	70	28	4	102	96.1%	5	24	1	30	96.7%
	2009	79	51	2	132	98.5%	78	34	2	114	98.2%	1	17		18	100.0%
	2010	79	67	1	147	99.3%	78	47		125	100.0%	1	20	1	22	95.5%
	2011	88	59	5	152	96.7%	84	38	5	127	96.1%	4	21		25	100.0%
	2012	102	64	6	172	96.5%	97	30	4	131	96.9%	5	34	2	41	95.1%
	2013	100	69	17	186	90.9%	98	54	15	167	91.0%	2	15	2	19	89.5%
	2014	45	56	64	165	61.2%	44	34	58	136	57.4%	1	22	6	29	79.3%
2005~2014	814	599	124	1,537	91.9%	790	383	112	1,285	91.3%	24	216	12	252	95.2%	
大腸 (結腸 +直腸)	2005	55	29	8	92	91.3%	54	26	8	88	90.9%	1	3		4	100.0%
	2006	86	41	4	131	96.9%	84	34	4	122	96.7%	2	7		9	100.0%
	2007	72	30	2	104	98.1%	71	25		96	100.0%	1	5	2	8	75.0%
	2008	77	25	2	104	98.1%	76	18	2	96	97.9%	1	7		8	100.0%
	2009	61	34	2	97	97.9%	60	30	2	92	97.8%	1	4		5	100.0%
	2010	75	34	1	110	99.1%	73	32	1	106	99.1%	2	2		4	100.0%
	2011	67	39	5	111	95.5%	65	30	5	100	95.0%	2	9		11	100.0%
	2012	84	41	7	132	94.7%	80	33	6	119	95.0%	4	8	1	13	92.3%
	2013	80	33	13	126	89.7%	78	26	13	117	88.9%	2	7		9	100.0%
	2014	43	40	43	126	65.9%	43	28	42	113	62.8%		12	1	13	92.3%
2005~2014	700	346	87	1,133	92.3%	684	282	83	1,049	92.1%	16	64	4	84	95.2%	
肝 (肝内 胆管 除く)	2005	2	20		22	100.0%	1	3		4	100.0%	1	17		18	100.0%
	2006	11	11	1	23	95.7%	7	2		9	100.0%	4	9	1	14	92.9%
	2007	6	22		28	100.0%	4	6		10	100.0%	2	16		18	100.0%
	2008	8	15	1	24	95.8%	7		1	8	87.5%	1	15		16	100.0%
	2009	8	13	1	22	95.5%	7	8		15	100.0%	1	5	1	7	85.7%
	2010	14	21		35	100.0%	11	6		17	100.0%	3	15		18	100.0%
	2011	10	22		32	100.0%	6	8		14	100.0%	4	14		18	100.0%
	2012	14	18	1	33	97.0%	10	9	1	20	95.0%	4	9		13	100.0%
	2013	7	14	3	24	87.5%	5	4	1	10	90.0%	2	10	2	14	85.7%
	2014	4	14	5	23	78.3%	4	3	5	12	58.3%		11		11	100.0%
2005~2014	84	170	12	266	95.5%	62	49	8	119	93.3%	22	121	4	147	97.3%	
膵	2005	2	34	1	37	97.3%	2	9		11	100.0%		25	1	26	96.2%
	2006	1	28		29	100.0%	1	12		13	100.0%		16		16	100.0%
	2007	1	24		25	100.0%	1	8		9	100.0%		16		16	100.0%
	2008	2	37		39	100.0%	2	11		13	100.0%		26		26	100.0%
	2009	6	30	1	37	97.3%	6	13	1	20	95.0%		17		17	100.0%
	2010	2	39	1	42	97.6%	2	12		14	100.0%		27	1	28	96.4%
	2011	4	36		40	100.0%	4	17		21	100.0%		19		19	100.0%
	2012	6	34		40	100.0%	4	7		11	100.0%	2	27		29	100.0%
	2013	1	44		45	100.0%	1	12		13	100.0%		32		32	100.0%
	2014	4	39	2	45	95.6%	4	15	1	20	95.0%		24	1	25	96.0%
2005~2014	29	345	5	379	98.7%	27	116	2	145	98.6%	2	229	3	234	98.7%	

※ここで示しているのは、診断から5年経過後の予後の把握状況であり、「5年生存率」ではありません。

原発部位	全症例 (上皮内がんを除く)(初発のみ)	手術(原発巣切除)あり	手術(原発巣切除)なし (姑息・対症療法ありを含む)
全部位	<p>不明 6.4%</p> <p>死亡 43.9%</p> <p>生存 49.7%</p>	<p>不明 7.3%</p> <p>死亡 28.0%</p> <p>生存 64.7%</p>	<p>不明 4.9%</p> <p>死亡 71.1%</p> <p>生存 24.0%</p>
食道	<p>不明 5.2%</p> <p>死亡 69.2%</p> <p>生存 25.6%</p>	<p>不明 5.6%</p> <p>死亡 57.6%</p> <p>生存 36.8%</p>	<p>不明 4.8%</p> <p>死亡 80.8%</p> <p>生存 14.4%</p>
胃	<p>不明 8.1%</p> <p>死亡 39.0%</p> <p>生存 53.0%</p>	<p>不明 8.7%</p> <p>死亡 29.8%</p> <p>生存 61.5%</p>	<p>不明 4.8%</p> <p>死亡 85.7%</p> <p>生存 9.5%</p>
大腸 (結腸 + 直腸)	<p>不明 7.7%</p> <p>死亡 30.5%</p> <p>生存 61.8%</p>	<p>不明 7.9%</p> <p>死亡 26.9%</p> <p>生存 65.2%</p>	<p>不明 4.8%</p> <p>死亡 76.2%</p> <p>生存 19.0%</p>
肝 (肝内胆管除く)	<p>不明 4.5%</p> <p>死亡 63.9%</p> <p>生存 31.6%</p>	<p>不明 6.7%</p> <p>死亡 41.2%</p> <p>生存 52.1%</p>	<p>不明 2.7%</p> <p>死亡 82.3%</p> <p>生存 15.0%</p>
膵	<p>不明 1.3%</p> <p>死亡 91.0%</p> <p>生存 7.7%</p>	<p>不明 1.4%</p> <p>死亡 80.0%</p> <p>生存 18.6%</p>	<p>不明 1.3%</p> <p>死亡 97.9%</p> <p>生存 0.9%</p>

原発部位	5年予後 診断年	全症例 (上皮内がんを除く)(初発のみ)					手術(原発巣切除)あり					手術(原発巣切除)なし (姑息・対症療法ありを含む)					
		生存	死亡	不明	計	予後 判明率	生存	死亡	不明	計	予後 判明率	生存	死亡	不明	計	予後 判明率	
肺	2005	42	82	1	125	99.2%	34	20		54	100.0%	8	62	1	71	98.6%	
	2006	40	93	2	135	98.5%	34	20	2	56	96.4%	6	73		79	100.0%	
	2007	33	71		104	100.0%	30	13		43	100.0%	3	58		61	100.0%	
	2008	40	89	2	131	98.5%	36	13	1	50	98.0%	4	76	1	81	98.8%	
	2009	43	94	1	138	99.3%	40	16	1	57	98.2%	3	78		81	100.0%	
	2010	41	101	2	144	98.6%	35	16	1	52	98.1%	6	85	1	92	98.9%	
	2011	53	115	5	173	97.1%	42	22	2	66	97.0%	11	93	3	107	97.2%	
	2012	49	139	9	197	95.4%	39	16	4	59	93.2%	10	123	5	138	96.4%	
	2013	56	93	8	157	94.9%	49	22	4	75	94.7%	7	71	4	82	95.1%	
	2014	54	105	37	196	81.1%	45	20	28	93	69.9%	9	85	9	103	91.3%	
2005~2014	451	982	67	1,500	95.5%	384	178	43	605	92.9%	67	804	24	895	97.3%		
乳房	2005	38	8	3	49	93.9%	37	5	3	45	93.3%	1	3		4	100.0%	
	2006	45	3	1	49	98.0%	45	3	1	49	98.0%						
	2007	50	5		55	100.0%	50	5		55	100.0%						
	2008	54	3	1	58	98.3%	54	2	1	57	98.2%		1		1	100.0%	
	2009	36	7	1	44	97.7%	36	6	1	43	97.7%		1		1	100.0%	
	2010	46	4	2	52	96.2%	45	3	2	50	96.0%	1	1		2	100.0%	
	2011	44	14	6	64	90.6%	42	11	2	55	96.4%	2	3	4	9	55.6%	
	2012	47	9	1	57	98.2%	47	6		53	100.0%		3	1	4	75.0%	
	2013	35	3	5	43	88.4%	35	3	4	42	90.5%				1	1	0.0%
	2014	37	7	24	68	64.7%	34	3	18	55	67.3%	3	4	6	13	53.8%	
2005~2014	432	63	44	539	91.8%	425	47	32	504	93.7%	7	16	12	35	65.7%		
子宮頸部	2005	17	8	3	28	89.3%	14	2	2	18	88.9%	3	6	1	10	90.0%	
	2006	13	4		17	100.0%	11	1		12	100.0%	2	3		5	100.0%	
	2007	7	2		9	100.0%	4			4	100.0%	3	2		5	100.0%	
	2008	7	3		10	100.0%	5	2		7	100.0%	2	1		3	100.0%	
	2009	14	5		19	100.0%	11	2		13	100.0%	3	3		6	100.0%	
	2010	13	3		16	100.0%	12			12	100.0%	1	3		4	100.0%	
	2011	14	5		19	100.0%	12			12	100.0%	2	5		7	100.0%	
	2012	22	6	5	33	84.8%	16	3	4	23	82.6%	6	3	1	10	90.0%	
	2013	20	2	7	29	75.9%	17	2	5	24	79.2%	3		2	5	60.0%	
	2014	19	7	10	36	72.2%	17	4	9	30	70.0%	2	3		6	83.3%	
2005~2014	146	45	25	216	88.4%	119	16	20	155	87.1%	27	29	5	61	91.8%		
子宮体部	2005	16	1	1	18	94.4%	16	1	1	18	94.4%						
	2006	10	4	2	16	87.5%	10	2	1	13	92.3%		2	1	3	66.7%	
	2007	20	4		24	100.0%	20	3		23	100.0%		1		1	100.0%	
	2008	15	3		18	100.0%	15	2		17	100.0%		1		1	100.0%	
	2009	17	4	3	24	87.5%	17	4	3	24	87.5%						
	2010	21	3		24	100.0%	21	2		23	100.0%		1		1	100.0%	
	2011	21	3	1	25	96.0%	21	3	1	25	96.0%						
	2012	19	4	2	25	92.0%	19	3	2	24	91.7%		1		1	100.0%	
	2013	22	3	2	27	92.6%	22	3	2	27	92.6%						
	2014	15	6	7	28	75.0%	15	2	7	24	70.8%		4		4	100.0%	
2005~2014	176	35	18	229	92.1%	176	25	17	218	92.2%	0	10	1	11	90.9%		
前立腺	2005	42	11	1	54	98.1%	17	2		19	100.0%	25	9	1	35	97.1%	
	2006	45	17	1	63	98.4%	13			13	100.0%	32	17	1	50	98.0%	
	2007	43	11		54	100.0%	16	1		17	100.0%	27	10		37	100.0%	
	2008	61	14		75	100.0%	28	3		31	100.0%	33	11		44	100.0%	
	2009	57	8	1	66	98.5%	27	2		29	100.0%	30	6	1	37	97.3%	
	2010	58	12	2	72	97.2%	32	3		35	100.0%	26	9	2	37	94.6%	
	2011	68	19	6	93	93.5%	29			29	100.0%	39	19	6	64	90.6%	
	2012	56	10	8	74	89.2%	22	1		23	100.0%	34	9	8	51	84.3%	
	2013	48	8	9	65	86.2%	24			24	100.0%	24	8	9	41	78.0%	
	2014	32	9	24	65	63.1%	20	4	11	35	68.6%	12	5	13	30	56.7%	
2005~2014	510	119	52	681	92.4%	228	16	11	255	95.7%	282	103	41	426	90.4%		

※ここで示しているのは、診断から5年経過後の予後の把握状況であり、「5年生存率」ではありません。

原発部位	全症例 (上皮内がんを除く)(初発のみ)	手術(原発巣切除)あり	手術(原発巣切除)なし (姑息・対症療法ありを含む)
肺	<p>不明 4.5% 生存 30.1% 死亡 65.5%</p>	<p>不明 7.1% 生存 63.5% 死亡 29.4%</p>	<p>不明 2.7% 生存 7.5% 死亡 89.8%</p>
乳房	<p>不明 8.2% 生存 80.1% 死亡 11.7%</p>	<p>不明 6.3% 生存 84.3% 死亡 9.3%</p>	<p>不明 34.3% 生存 20.0% 死亡 45.7%</p>
子宮頸部	<p>不明 11.6% 生存 67.6% 死亡 20.8%</p>	<p>不明 12.9% 生存 76.8% 死亡 10.3%</p>	<p>不明 8.2% 生存 44.3% 死亡 47.5%</p>
子宮体部	<p>不明 7.9% 生存 76.9% 死亡 15.3%</p>	<p>不明 7.8% 生存 80.7% 死亡 11.5%</p>	<p>不明 9.1% 生存 0.0% 死亡 90.9%</p>
前立腺	<p>不明 7.6% 生存 74.9% 死亡 17.5%</p>	<p>不明 4.3% 生存 89.4% 死亡 6.3%</p>	<p>不明 9.6% 生存 66.2% 死亡 24.2%</p>